

***Eriogonum heermannii* Durand & Hilgard *var. clokeyi* Reveal (1976) CLOKEY BUCKWHEAT**

FAMILY: *Polygonaceae*, the buckwheat family.

STATUS: **Heritage Program SENSITIVE LIST, ranks: G5 T2 S2**

USFWS/ESA: none. **STATE OF NEVADA:** none. **BLM:** Special Status Species. **USFS:** Humboldt-Toiyabe NF Sensitive Species. **NNNPS:** watch list.

POPULATION CENSUS: 8 occurrences mapped; total estimated individuals 459+, total estimated area unknown.

TREND: unknown.

IMPACTS AND MAJOR THREATS: No summary available (see references).

INVENTORY EFFORT: Not yet systematically surveyed. Most recent entered survey 1987, average year of last survey 1968. **Years since last entered survey** (percent of mapped records at various survey ages): **11-20 yrs:** 12.5%; **21-30 yrs:** 12.5%; **31-50 yrs:** 62.5%; **51+ yrs or unknown:** 12.5%.

LAND MANAGEMENT in decreasing predominance: Humboldt-Toiyabe National Forest, U. S. Bureau of Land Management, U. S. Department of Energy, U. S. Fish and Wildlife Service, U. S. Department of Defense, private lands (?).

RANGE: Clark and Nye counties, Nevada. **Nevada endemic.** Maximum **range dimension 71.9 km (44.7 mi)** excluding most disjunct record. **Type specimen** collected in Clark County.

ELEVATIONS RECORDED: 4000-6000 feet (1219-1829 meters).

HABITAT: Carbonate outcrops, talus, scree, and gravelly washes and banks in the creosote-bursage, shadscale, and blackbrush zones.

PHENOLOGY: flowering late-spring to summer. Range of most frequent **survey months:** June-September.

LIFE-FORM AND HABIT: shrub.

DESCRIPTION: not available (see references).

PHOTOGRAPHS: none known.

ILLUSTRATIONS: none known.

OTHER GENERAL REFERENCES (listed separately): Clokey (1951), Reveal (1976), Reveal (1985).
